arXiv:1409.5156v5 [math.FA] 24 Mar 2016

THE WEIGHTED MEAN MATRIX WITH WEIGHT SEQUENCE $w_n = 2n + 1$ IS A HYPONORMAL OPERATOR ON ℓ^2

H. C. RHALY JR.

In memory of Russell Aubrey Stokes (1922-2014)

Abstract. A weighted mean matrix whose weight sequence is linear with positive coefficients is shown to be a posinormal operator on ℓ^2. This operator is also shown to be coposinormal, so it and its adjoint have the same null space and the same range. The posinormality result leads to a proof that the weighted mean matrix generated by the sequence of odd positive integers is hyponormal, as well as a conjecture regarding the more general case.

1. Introduction

If $\mathcal{B}(H)$ is the set of all bounded linear operators on a Hilbert space H, then the operator $A \in \mathcal{B}(H)$ is *hyponormal* if

$$< (A^*A - AA^*)f, f > \geq 0$$

for all $f \in H$, and A is said to be *posinormal* if

$$AA^* = A^*PA$$

for some positive operator $P \in \mathcal{B}(H)$, called the *interrupter*; A is *coposinormal* if A^* is posinormal. The following proposition (see [5]) contains some key facts about posinormal operators.

Proposition 1.1. *If $A \in \mathcal{B}(H)$, then the following are equivalent.*

(a) *There exists a nonnegative $P \in \mathcal{B}(H)$ such that $AA^* = A^*PA$ (i.e., A is posinormal).*
(b) *There exists a nonnegative $P \in \mathcal{B}(H)$ such that $AA^* \leq A^*PA$.*
(c) *There exists a nonnegative $\alpha \in \mathbb{R}$ such that $AA^* \leq \alpha^2 A^*A$.*
(d) *$Ran(A) \subseteq Ran(A^*)$, where $Ran(A) = \{g \in H : g = Af$ for some $f \in H\}$.*

2010 *Mathematics Subject Classification.* 47B20, 47B37.
Key words and phrases. hyponormal operator, posinormal operator, weighted mean matrix.

(e) *There exists a* $B \in \mathcal{B}(H)$ *such that* $A = A^*B$.
Moreover, each of the above assertions implies the following one.

(f) $Ker(A) \subseteq Ker(A^*)$, *where* $Ker(A) = \{f \in H : Af = 0\}$.
Furthermore, if $Ran(A)$ *is closed, then these six assertions are all equivalent.*

Proof. See [5]. □

Note that A is hyponormal if part (c) of the proposition is satisfied for $\alpha = 1$, so hyponormal operators are necessarily posinormal.

A lower triangular infinite matrix $M = [m_{ij}] \in \mathcal{B}(\ell^2)$ is *factorable* if its entries are of the form

$$m_{ij} = \begin{cases} a_i c_j & if \quad j \leq i \\ 0 & if \quad j > i \end{cases}$$

where a_i depends only on i and c_j depends only on j. A *weighted mean matrix* is a lower triangular matrix with entries w_j/W_i, where $\{w_j\}$ is a nonnegative sequence with $w_0 > 0$, and $W_i = \sum_{j=0}^{i} w_j$. A weighted mean matrix is factorable, with $a_i = 1/W_i$ and $c_j = w_j$ for all i,j. In recent decades these operators, acting on various sequence spaces, have been studied by many authors in a number of papers, including [1], [2], [3], [4], [7], [12],and [13]. Hyponormality on ℓ^2 has been demonstrated in three special cases, namely, for the operators generated by $w_j = j + 1$ (see [10]), $w_j = (j + 1)^2$ (see [11]), and $w_j = 2 - 1/2^j$ (see [9]); the first and third examples were handled without software assistance, in contrast with what will be the case in this paper.

Initially under consideration here will be the weighted mean matrix M associated with the general linear weight sequence $w_n = an + b$ where a, $b > 0$. The first aim is to show that this operator is both posinormal and coposinormal. Posinormality will be useful for achieving the two main objectives of this paper, (1) showing that the matrix associated with the sequence $w_n = 2n + 1$ is a hyponormal operator on ℓ^2 and (2) propounding a conjecture for hyponormality of the more general case.

2. Posinormality and hyponormality

Our first goal is to demonstrate that a weighted mean matrix M whose weight sequence is linear with positive coefficients is a posinormal operator on ℓ^2. The following theorem from [8] provides a tool for that purpose.

Theorem 2.1. *Suppose* $M = [a_i c_j]$ *is a lower triangular factorable matrix that acts as a bounded operator on* ℓ^2 *and that the following conditions are satisfied:*

(a) *both* $\{a_n\}$ *and* $\{a_n/c_n\}$ *are positive decreasing sequences that converge to 0, and*
(b) *the matrix* B *defined by* $B = [b_{ij}]$ *by*

$$b_{ij} = \begin{cases} c_i(\frac{1}{c_j} - \frac{1}{c_{j+1}}\frac{a_{j+1}}{a_j}) & if \quad i \leq j; \\ -\frac{a_{j+1}}{a_j} & if \quad i = j+1; \\ 0 & if \quad i > j+1. \end{cases}$$

is a bounded operator on ℓ^2.

Then M *is posinormal.*

Proof. See [8]. □

Theorem 2.2. *The weighted mean matrix* M *associated with the weight sequence* $w_n = an + b$ *with* a, $b > 0$ *is a posinormal operator on* ℓ^2.

Proof. It suffices to consider sequences of the form $w_n = kn+1$ with $k > 0$. It is clear that condition (a) of Theorem 2.1 is satisfied, so we focus our attention now on condition (b). Since $c_n = kn + 1$ and $a_n = \frac{2}{(n+1)(kn+2)}$, the matrix $B = [b_{ij}]$ entries become

$$b_{ij} = \begin{cases} (ki+1) \cdot \frac{3k^2j^2+(5k^2+6k)j+2k^2+6k+2}{(j+2)(kj+1)(kj+k+1)(kj+k+2)} & if \quad i \leq j; \\ -\frac{(j+1)(kj+2)}{(j+2)(kj+k+2)} & if \quad i = j+1; \\ 0 & if \quad i > j+1. \end{cases}$$

Note that for $i \leq j$, $0 < b_{ij} \leq 3/(j+2)$ for all j, so the upper triangular part T of matrix B is a bounded operator on ℓ^2. If W is the weighted shift with weight sequence $\{\frac{(n+1)(kn+2)}{(n+2)(kn+k+2)} : n \geq 0\}$, then $B = T - W$, so $B \in \mathcal{B}(\ell^2)$. It now follows from Theorem 2.1 that M is posinormal. □

We note that the range of M contains all the e_n's from the standard orthonormal basis for ℓ^2. From [8] we know that the entries of the interrupter $P = [p_{ij}] = B^*B$ are given by

$$p_{ij} = \begin{cases} \frac{c_j^2c_{j+1}^2a_{j+1}^2+(\sum_{\ell=0}^{j}c_\ell^2)(c_{j+1}a_j-c_ja_{j+1})^2}{c_j^2c_{j+1}^2a_j^2} & if \quad i = j; \\ \frac{(c_ia_{i+1}-c_{i+1}a_i)[c_j(\sum_{\ell=0}^{j+1}c_\ell^2)a_{j+1}-c_{j+1}(\sum_{\ell=0}^{j}c_\ell^2)a_j]}{c_ic_{i+1}c_jc_{j+1}a_ia_j} & if \quad i > j; \\ \frac{(c_ja_{j+1}-c_{j+1}a_j)[c_i(\sum_{\ell=0}^{i+1}c_\ell^2)a_{i+1}-c_{i+1}(\sum_{\ell=0}^{i}c_\ell^2)a_i]}{c_ic_{i+1}c_jc_{j+1}a_ia_j} & if \quad i < j. \end{cases}$$

For M to be hyponormal, it must be true that

$$\langle (M^*M - MM^*)f, f\rangle = \langle (M^*M - M^*PM)f, f\rangle = \langle (I-P)Mf, Mf\rangle \geq 0$$

for all f in ℓ^2. Consequently, we can conclude that M will be hyponormal when $Q :\equiv I - P \geq 0$. Again using $c_n = kn + 1$ and $a_n = \frac{2}{(n+1)(kn+2)}$, we determine that the entries of $Q = [q_{ij}]$ are given by $q_{ij} =$

$$\begin{cases} \frac{6k^6j^6+(21k^6+42k^5)j^5+(25k^6+126k^5+114k^4)j^4+(9k^6+126k^5+266k^4+168k^3)j^3}{6(j+2)(kj+1)^2(kj+k+1)^2(kj+k+2)^2} & \\ \quad +\frac{(-3k^6+42k^5+190k^4+264k^3+160k^2)j^2+(-2k^6+38k^4+96k^3+148k^2+96k)j}{6(j+2)(kj+1)^2(kj+k+1)^2(kj+k+2)^2} & \\ \quad +\frac{12k^2+48k+24}{6(j+2)(kj+1)^2(kj+k+1)^2(kj+k+2)^2} & if \quad i = j; \\ -\frac{1}{6}\cdot\frac{3k^2i^2+(5k^2+6k)i+(2k^2+6k+2)}{(i+2)(ki+1)(ki+k+1)(ki+k+2)}\cdot\frac{(j+1)(k^4j^2+(k^4+4k^2)j+6k)}{(kj+1)(kj+k+1)(kj+k+2)} & if \quad i > j; \\ -\frac{1}{6}\cdot\frac{3k^2j^2+(5k^2+6k)j+(2k^2+6k+2)}{(j+2)(kj+1)(kj+k+1)(kj+k+2)}\cdot\frac{(i+1)(k^4i^2+(k^4+4k^2)i+6k)}{(ki+1)(ki+k+1)(ki+k+2)} & if \quad i < j. \end{cases} \tag{2.1}$$

When $k = 1$, Q assumes the form studied in [10]. When $k \neq 1$, the key lemma in [10] is not satisfied, so a different approach is required here. Continuing toward hyponormality in the general case has heretofore not been fruitful, so we focus our attention now on the case $k = 2$.

Theorem 2.3. *The weighted mean matrix M associated with the weight sequence $w_n = 2n + 1$ is a hyponormal operator on ℓ^2.*

Proof. Using $k = 2$, we find that the entries of $Q = [q_{ij}]$, in irreducible form, are

$$q_{ij} = \begin{cases} \frac{12j^4+60j^3+104j^2+66j+7}{3(j+2)^3(2j+1)(2j+3)} & if \quad i = j; \\ -\frac{1}{3}\cdot\frac{6i^2+16i+11}{(i+2)^2(2i+1)(2i+3)}\cdot\frac{j+1}{j+2} & if \quad i > j; \\ -\frac{1}{3}\cdot\frac{6j^2+16j+11}{(j+2)^2(2j+1)(2j+3)}\cdot\frac{i+1}{i+2} & if \quad i < j. \end{cases}$$

In order to show that Q is positive, it suffices to show that Q_N, the N^{th} finite section of Q (involving rows $i = 0, 1, 2, ..., N$ and columns $j = 0, 1, 2, ..., N$), has positive determinant for each positive integer N. For columns $j = 0, 1, ...,$ N-1, we multiply the $(j + 1)^{st}$ column of Q_N by $z_j :\equiv \frac{(j+1)(j+3)}{(j+2)^2}$ and subtract from the j^{th} column. Call the new matrix Q'_N. Then we work with the rows of Q'_N. For $i = 0, 1, ..., N - 1$, we multiply the $(i + 1)^{st}$ row of Q'_N by z_i and subtract from the i^{th} row. This leads to the tridiagonal form

$$Y_N :\equiv \begin{pmatrix} d_0 & s_0 & 0 & \dots & 0 & 0 \\ s_0 & d_1 & s_1 & \dots & 0 & 0 \\ 0 & s_1 & d_2 & \dots & . & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & . & \dots & d_{N-1} & s_{N-1} \\ 0 & 0 & 0 & \dots & s_{N-1} & d_N \end{pmatrix},$$

where

$$d_n = q_{nn} - z_n q_{n,n+1} - z_n(q_{n+1,n} - z_n q_{n+1,n+1}) = q_{nn} - 2z_n q_{n,n+1} + z_n^2 q_{n+1,n+1}$$
$$= \frac{16n^7 + 200n^6 + 1024n^5 + 2768n^4 + 4226n^3 + 3576n^2 + 1481n + 197}{(n+2)^4(n+3)(2n+1)(2n+3)(2n+5)}$$

and

$$s_n = q_{n+1,n} - z_n q_{n+1,n+1} = -\frac{(n+1)(2n^2+8n+7)}{(n+2)^2(n+3)(2n+3)} \text{ when } 0 \le n \le N-1;$$

and lastly,

$$d_N = q_{NN} = \frac{12N^4 + 60N^3 + 104N^2 + 66n + 7}{3(N+2)^3(2N+1)(2N+3)}.$$

Note that $\det Y_N = \det Q_N^{'} = \det Q_N$. Next we transform Y_N into a triangular matrix with the same determinant. The new matrix has diagonal entries δ_n which are given by the recursion formula: $\delta_0 = d_0$, $\delta_n = d_n - s_{n-1}^2/\delta_{n-1}$ $(1 \le n \le N)$. An induction argument shows that

$$\delta_n > \frac{4n+10}{4n^2+20n+37}$$

for $0 \le n \le N-1$. Since d_N departs from the pattern set by the earlier d_n's, δ_N needs to be handled separately:

$$\delta_N \ge \frac{24N^8 + 140N^7 + 432N^6 + 1160N^5 + 2234N^4 + 2297N^3 + 1070N^2 + 216N + 14}{6(N+1)^4(N+2)^3(2N+1)^2(2N+3)}.$$

Therefore $\det Q_N = \prod_{n=0}^{N} \delta_n > 0$, so the proof is complete. □

The verification of the induction step above reduces – using the SageMath software system [14] – to the observation that $96n^{10} + 4944n^9 + 54624n^8 + 282288n^7 + 824294n^6 + 1447767n^5 + 1531563n^4 + 927504n^3 + 285132n^2 + 35022n + 178 \ge 0$ for all $n \ge 1$. The reduction was accomplished when SageMath executed the following command.

```
expand((16*n^7+200*n^6+1024*n^5+2768*n^4+4226*n^3+3576*n^2+
1481*n+197)*2*(n+1)^4*(2*n+1)*(4*n^2+20*n+37)-n^2*(2*n^2+4*
n+1)^2*(4*n^2+12*n+21)*(n+2)^2*(n+3)*(2*n+5)*(4*n^2+20*n+37)-
(4*n+10)*2*(n+1)^4*(n+2)^4*(n+3)*(2*n+1)^2*(2*n+3)*(2*n+5))
```

3. Coposinormality

We now proceed to show that a weighted mean operator whose weight sequence is linear with positive coefficients has a posinormal adjoint. The next theorem provides a tool for doing that.

Theorem 3.1. *Suppose $M = [a_i c_j]$ is a lower triangular factorable matrix that acts as a bounded operator on ℓ^2 and that the following conditions are satisfied:*

(a) *both $\{a_n\}$ and $\{a_n/c_n\}$ are positive decreasing sequences that converge to 0, and*
(b) *the matrix $Z = [z_{ij}]$ defined by*

$$z_{ij} = \begin{cases} \frac{a_i}{a_j} & if \quad j = 0; \\ a_i(\frac{1}{a_j} - \frac{c_{j-1}}{c_j}\frac{1}{a_{j-1}}) & if \quad 0 < j \leq i; \\ -\frac{c_{j-1}}{c_j} & if \quad j = i+1; \\ 0 & if \quad j > i+1. \end{cases}$$

is a bounded operator on ℓ^2,

Then M^* is posinormal.

Proof. See [8]. □

Theorem 3.2. *The weighted mean matrix M associated with the weight sequence $w_n = an + b$ with a, $b > 0$ is a coposinormal operator on ℓ^2.*

Proof. Again it suffices to consider sequences of the form $w_n = kn + 1$ with $k > 0$. It is clear that condition (a) of Theorem 3.1 is satisfied, so we focus our attention now on condition (b). When $c_n = kn + 1$ and $a_n = \frac{2}{(n+1)(kn+2)}$, the matrix $Z = [z_{ij}]$ becomes

$$z_{ij} = \begin{cases} \frac{2}{(i+1)(ki+2)} & if \quad j = 0; \\ \frac{1}{(i+1)(ki+2)} \cdot \frac{3k^2j^2+(6k-k^2)j+2}{kj+1} & if \quad 0 < j \leq i; \\ -\frac{kj-k+1}{kj+1} & if \quad j = i+1; \\ 0 & if \quad j > i+1. \end{cases}$$

Note that for $j \leq i$, $0 < z_{ij} \leq 3/(i+1)$ for all i, so the lower triangular part T of matrix Z is a bounded operator on ℓ^2. If W is the weighted shift with weight sequence $\{(kn+1)/(kn+k+1) : n \geq 0\}$, then $Z = T - W^*$, so $Z \in B(\ell^2)$. It now follows from Theorem 3.1 that M^* is posinormal. □

Corollary 3.1. *If M is the weighted mean operator on ℓ^2 determined by the sequence $w_n = an + b$ with a, $b > 0$, then M^j is both posinormal and coposinormal for each positive integer j.*

Proof. This follows from Theorems 2.2 and 3.2, together with [6, Corollary 2.2(b)]. □

Theorem 3.3. *If M is the weighted mean operator on ℓ^2 determined by the sequence $w_n = an + b$ with a, $b > 0$, then both M and M^* are injective and have dense range with*

$$Ran(M) = Ran(M^*).$$

Proof. Since M is both posinormal and coposinormal, it follows from Proposition 1.1 that $Ker(M) = Ker(M^*)$ and $Ran(M) = Ran(M^*)$. It is easy to see that $Ker(M) = \{0\}$. Consequently, both M and M^* are one-to-one, and both have dense range. □

4. Concluding Remarks

After seeing the hyponormality result for $w_n = 2n+1$, one might reasonably wonder whether the calculations for the weighted mean matrix for weight sequence $w_n = 3n+1$ are significantly more complex, and the answer is – Yes. To illustrate, we observe that for $w_n = 2n + 1$, the irreducible form of q_{nn} has a degree 4 numerator and a degree 5 denominator (see the proof of Theorem 2.3). In contrast, for $w_n = 3n + 1$, the irreducible form of q_{nn} has a degree 6 numerator and a degree 7 denominator – as well as larger coefficients for all the terms, so the hyponormality calculations for that weight sequence are significantly more complicated than what was presented here. Nevertheless, we offer the following for the reader to consider:

- Successful calculations have been run for $w_n = kn + 1$ when $k = 1/2$, $4/3$, $3/2$, 3, and 4, besides $k = 2$ (for $k = 1$, see [10]).
- The first four finite sections of Q, whose general entries are displayed in 2.1, have positive determinants for all $k > 0$ (see Appendix).

Consequently, we close with the following conjecture.

Conjecture 4.1. *If M is a weighted mean matrix whose weight sequence is linear with positive coefficients, then M is a hyponormal operator on ℓ^2.*

5. Appendix

The expressions below refer to the second bulleted item immediately preceding Conjecture 4.1. The determinant of the first finite section of Q is

$$\det(Q_0) = \frac{k^2 + 4k + 2}{(k+1)^2(k+2)^2},$$

the determinant of the second is

$$\det(Q_1) = \frac{7k^6 + 56k^5 + 114k^4 + 110k^3 + 76k^2 + 36k + 6}{9(k+1)^4(k+2)^2(2k+1)^2},$$

the determinant of the third is

$$\det(Q_2) = \frac{405k^{10} + 4320k^9 + 13959k^8 + 22968k^7 + 24214k^6 + 18464k^5}{36(k+1)^4(k+2)^2(2k+1)^2(3k+1)^2(3k+2)^2} + \frac{10527k^4 + 4496k^3 + 1423k^2 + 288k + 24}{36(k+1)^4(k+2)^2(2k+1)^2(3k+1)^2(3k+2)^2},$$

and the determinant of the fourth is

$$\begin{aligned}\det(Q_3) = &\frac{101088k^{14} + 1280448k^{13} + 5308128k^{12} + 11654028k^{11}}{900(k+1)^4(k+2)^2(2k+1)^4(3k+1)^2(3k+2)^2(4k+1)^2} \\ &+ \frac{16473861k^{10} + 16665540k^9 + 12793267k^8 + 7701000k^7}{900(k+1)^4(k+2)^2(2k+1)^4(3k+1)^2(3k+2)^2(4k+1)^2} \\ &+ \frac{3719530k^6 + 1460772k^5 + 463575k^4 + 115268k^3}{900(k+1)^4(k+2)^2(2k+1)^4(3k+1)^2(3k+2)^2(4k+1)^2} \\ &+ \frac{20975k^2 + 2400k + 120}{900(k+1)^4(k+2)^2(2k+1)^4(3k+1)^2(3k+2)^2(4k+1)^2}.\end{aligned}$$

These calculations were assisted by [14]. Clearly each of these determinants is positive for all $k > 0$.

References

[1] D. Borwein and X. Gao, *Generalized Hausdorff and weighted mean matrices as operators on l_p*, J. Math. Anal. Appl. **178** (1993), no. 2, 517–528.

[2] P. Gao, *On l^p norms of weighted mean matrices*, Math. Z. **264** (2010), no. 4, 829–848.

[3] P. Gao, *A note on l^p norms of weighted mean matrices*, J. Inequal. Appl. **2012**, 2012:110, 7 pp.

[4] C. E. Harrell, *Riesz matrices that are also Hausdorff matrices*, Proc. Amer. Math. Soc. **22** 1969 303–304.

[5] C. S. Kubrusly and B. P. Duggal, *On posinormal operators*, Adv. Math. Sci. Appl. **120** (2007), no. 1, 131-147.

[6] C. S. Kubrusly, P. C. M. Vieira, and J. Zanni, *Powers of posinormal operators*, Oper. Matrices, **10** (2016), no. 1, 15-27.

[7] E. Pazouki and B. Yousefi, *The spectra and eigenvectors for the weighted mean matrix operator*, Arab J. Math. Sci. **21** (2015), no. 2, 179–193.

[8] H. C. Rhaly Jr., *Posinormal factorable matrices whose interrupter is diagonal*, Mathematica (Cluj) **53 (76)** (2011), no. 2, 181-188.

[9] H. C. Rhaly Jr. and B. E. Rhoades, *Conditions for factorable matrices to be hyponormal and dominant*, Sib. Èlektron. Mat. Izv. **9** (2012), 261–265.

[10] H. C. Rhaly Jr. and B. E. Rhoades, *The weighted mean operator on ℓ^2 with the weight sequence $w_n = n + 1$ is hyponormal*, New Zealand J. Math. **44** (2014), 103-106.

[11] H. C. Rhaly Jr. and B. E. Rhoades, *The weighted mean operator on ℓ^2 with weight sequence $w_n = (n + 1)^p$ is hyponormal for $p = 2$*, Stud. Univ. Babeş-Bolyai Math. **59** (2014), no. 3, 365–368

[12] B. E. Rhoades, *The spectra of weighted mean operators on* bv_0, J. Austral. Math. Soc. Ser. A **52** (1992), no. 2, 242–250.

[13] B. E. Rhoades, *The equivalence of Cesàro and weighted mean matrices on various sequence spaces*, Indian J. Math. **36** (1994), no. 1, 1–6.
[14] W. A. Stein et al., Sage Mathematics Software (Version 6.2), The Sage Development Team, 2014, http://www.sagemath.org.

1081 Buckley Drive, Jackson, MS 39206, U.S.A.
E-mail address: rhaly@member.ams.org